

PRACTICE GUIDELINES FOR OBSTETRIC ANESTHESIA
*An Updated Report by the American Society of Anesthesiologists Task Force on
Obstetric Anesthesia**
(Last amended on October 18, 2006)

PRACTICE guidelines are systematically developed recommendations that assist the practitioner and patient in making decisions about health care. These recommendations may be adopted, modified, or rejected according to clinical needs and constraints and are not intended to replace local institutional policies. In addition, practice guidelines are not intended as standards or absolute requirements, and their use cannot guarantee any specific outcome. Practice guidelines are subject to revision as warranted by the evolution of medical knowledge, technology, and practice. They provide basic recommendations that are supported by a synthesis and analysis of the current literature, expert opinion, open forum commentary, and clinical feasibility data.

This update includes data published since the "Practice Guidelines for Obstetrical Anesthesia" were adopted by the American Society of Anesthesiologists in 1998; it also includes data and recommendations for a wider range of techniques than was previously addressed.

Methodology

A. Definition of Perioperative Obstetric Anesthesia

For the purposes of these Guidelines, obstetric anesthesia refers to peripartum anesthetic and analgesic activities performed during labor and vaginal delivery, cesarean delivery, removal of retained placenta, and postpartum tubal ligation.

* Developed by the American Society of Anesthesiologists Task Force on Obstetric Anesthesia: Joy L. Hawkins, M.D. (Chair), Denver, CO; James F. Arens, M.D., Houston, TX; Brenda A Bucklin, M.D., Denver, CO; Richard T. Connis, Ph.D., Woodinville, WA; Patricia A. Dailey, M.D., Hillsborough, CA; David R. Gambling, M.B.B.S., San Diego, CA; David G. Nickinovich, Ph.D., Bellevue, WA; Linda S. Polley, M.D., Ann Arbor, MI; Lawrence C. Tsen, M.D., Boston, MA; David J. Wlody, M.D., Brooklyn, NY; and Kathryn J. Zuspan, M.D., Stillwater, MN. Supported by the American Society of Anesthesiologists under the direction of James F. Arens, M.D., Chair, Committee on Practice Parameters. Approved by the House of Delegates on October 18, 2006. A list of the references used to develop these Guidelines is available by writing to the American Society of Anesthesiologists. Address reprint requests to the American Society of Anesthesiologists: 520 N. Northwest Highway, Park Ridge, Illinois 60068-2573.

B. Purposes of the Guidelines

The purposes of these Guidelines are to enhance the quality of anesthetic care for obstetric patients, improve patient safety by reducing the incidence and severity of anesthesia-related complications, and increase patient satisfaction.

C. Focus

These Guidelines focus on the anesthetic management of pregnant patients during labor, non-operative delivery, operative delivery, and selected aspects of postpartum care and analgesia (i.e., neuraxial opioids for postpartum analgesia after neuraxial anesthesia for cesarean delivery). The intended patient population includes, but is not limited to intrapartum and postpartum patients with uncomplicated pregnancies or with common obstetric problems. The Guidelines do not apply to patients undergoing surgery during pregnancy, gynecological patients or parturients with chronic medical disease (e.g., severe cardiac, renal or neurological disease). In addition, these Guidelines do not address: (1) postpartum analgesia for vaginal delivery, (2) analgesia after tubal ligation, or (3) postoperative analgesia after general anesthesia for cesarean delivery.

D. Application

These Guidelines are intended for use by anesthesiologists. They also may serve as a resource for other anesthesia providers and health care professionals who advise or care for patients who will receive anesthetic care during labor, delivery and the immediate postpartum period.

E. Task Force Members and Consultants

The American Society of Anesthesiologists (ASA) appointed a Task Force of eleven members to (1) review the published evidence, (2) obtain the opinion of a panel of consultants including anesthesiologists and non-anesthesiologist physicians concerned with obstetric anesthesia and analgesia, and (3) obtain opinions from practitioners likely to be affected by the Guidelines. The Task Force included anesthesiologists in both private and academic practices from various geographic areas

of the United States and two consulting methodologists from the ASA Committee on Practice Parameters.

The Task Force developed the Guidelines by means of a seven-step process. First, they reached consensus on the criteria for evidence. Second, original published research studies from peer-reviewed journals relevant to obstetric anesthesia were reviewed. Third, the panel of expert consultants was asked to: (a) participate in opinion surveys on the effectiveness of various peripartum management strategies and (b) review and comment on a draft of the Guidelines developed by the Task Force. Fourth, opinions about the Guideline recommendations were solicited from active members of the ASA who provide obstetric anesthesia. Fifth, the Task Force held open forums at two major national meetings[†] to solicit input on its draft recommendations. Sixth, the consultants were surveyed to assess their opinions on the feasibility of implementing the Guidelines. Seventh, all available information was used to build consensus within the Task Force to finalize the Guidelines.

F. Availability and Strength of Evidence

Preparation of these Guidelines followed a rigorous methodological process. To convey the findings in a concise and easy to understand fashion, these Guidelines employ several descriptive terms. When sufficient numbers of studies are available for evaluation, the following terms describe the strength of the findings.

Support: Meta-analysis of a sufficient number of randomized controlled trials[‡] indicates a statistically significant relationship ($p < 0.01$) between a clinical intervention and a clinical outcome.

Suggest: Information from case reports and observational studies permits inference of a relationship between an intervention and an outcome. A meta-analytic assessment of this type of qualitative or descriptive information is not conducted.

[†] International Anesthesia Research Society, 80th Clinical and Scientific Congress, March 25, 2006 in San Francisco, CA; and Society of Obstetric Anesthesia and Perinatology 38th Annual Meeting, April 29, 2006 in Hollywood, FL.

[‡] A prospective nonrandomized controlled trial may be included in a meta-analysis under certain circumstances if specific statistical criteria are met.

Equivocal: Either a meta-analysis has not found significant differences among groups or conditions, or there is insufficient quantitative information to conduct a meta-analysis and information collected from case reports and observational studies does *not* permit inference of a relationship between an intervention and an outcome.

The *lack* of scientific evidence in the literature is described by the following terms.

Silent: No identified studies address the specified relationship between an intervention and outcome.

Insufficient: There are too few published studies to investigate a relationship between an intervention and outcome.

Inadequate: The available studies cannot be used to assess the relationship between an intervention and an outcome. These studies either do not meet the criteria for content as defined in the “Focus” of these Guidelines, or do not permit a clear causal interpretation of findings due to methodological concerns.

Formal survey information is collected from consultants and members of the ASA. The following terms describe survey responses for any specified issue. Responses are solicited on a 5-point scale; ranging from 1 (strongly disagree) to 5 (strongly agree) with a score of 3 being equivocal. Survey responses are summarized based on median values as follows:

- Strongly Agree:* Median score of 5 (At least 50% of the responses are 5)
- Agree:* Median score of 4 (At least 50% of the responses are 4 or 4 and 5)
- Equivocal:* Median score of 3 (At least 50% of the responses are 3, or no other response category or combination of similar categories contain at least 50% of the responses)
- Disagree:* Median score of 2 (At least 50% of responses are 2 or 1 and 2)
- Strongly Disagree:* Median score of 1 (At least 50% of responses are 1)

Guidelines:

I. Perianesthetic Evaluation

History and Physical Examination. Although comparative studies are insufficient to evaluate the peripartum impact of conducting a focused history (e.g., reviewing medical records) or a physical examination, the literature reports certain patient or clinical characteristics that may be associated with obstetric complications. These characteristics include, but are not limited to, preeclampsia, pregnancy-related hypertensive disorders, HELLP syndrome, obesity and diabetes.

The consultants and ASA members both strongly agree that a directed history and physical examination, as well as communication between anesthetic and obstetric providers, reduces maternal, fetal and neonatal complications.

Recommendations. The anesthesiologist should conduct a focused history and physical examination before providing anesthesia care. This should include, but is not limited to, a maternal health and anesthetic history, a relevant obstetric history, a baseline blood pressure measurement, and an airway, heart, and lung examination, consistent with the ASA “Practice Advisory for Preanesthesia Evaluation.”[§]. When a neuraxial anesthetic is planned or placed, the patient’s back should be examined.

Recognition of significant anesthetic or obstetric risk factors should encourage consultation between the obstetrician and the anesthesiologist. A communication system should be in place to encourage early and ongoing contact between obstetric providers, anesthesiologists, and other members of the multidisciplinary team.

Intrapartum Platelet Count. The literature is insufficient to assess whether a routine platelet count can predict anesthesia-related complications in uncomplicated parturients. The literature suggests that a platelet count is clinically useful for parturients with suspected pregnancy-related

[§] American Society of Anesthesiologists Task Force on Preanesthesia Evaluation: Practice advisory for preanesthesia evaluation. *Anesthesiology* 2002; 96:485-496.

hypertensive disorders, such as preeclampsia or HELLP syndrome, and for other disorders associated with coagulopathy.

The ASA members are equivocal, but the consultants agree that obtaining a routine intrapartum platelet count does *not* reduce maternal anesthetic complications. Both the consultants and ASA members agree that, for patients with suspected preeclampsia, a platelet count reduces maternal anesthetic complications. The consultants strongly agree and the ASA members agree that a platelet count reduces maternal anesthetic complications for patients with suspected coagulopathy.

Recommendations. A specific platelet count predictive of neuraxial anesthetic complications has not been determined. The anesthesiologist's decision to order or require a platelet count should be individualized and based on a patient's history, physical examination and clinical signs. A routine platelet count is not necessary in the healthy parturient.

Blood Type and Screen. The literature is insufficient to determine whether obtaining a blood type and screen is associated with fewer maternal anesthetic complications. In addition, the literature is insufficient to determine whether a blood cross-match is necessary for healthy and uncomplicated parturients. The consultants and ASA members agree that an intrapartum blood sample should be sent to the blood bank for all parturients.

Recommendations. A routine blood cross-match is not necessary for healthy and uncomplicated parturients for vaginal or operative delivery. The decision whether to order or require a blood type and screen, or cross-match, should be based on maternal history, anticipated hemorrhagic complications (*e.g.*, placenta accreta in a patient with placenta previa and previous uterine surgery), and local institutional policies.

Perianesthetic Recording of the Fetal Heart Rate. The literature suggests that anesthetic and analgesic agents may influence the fetal heart rate pattern. There is insufficient literature to demonstrate that perianesthetic recording of the fetal heart rate prevents fetal or neonatal

complications. Both the consultants and ASA members agree, however, that perianesthetic recording of the fetal heart rate reduces fetal and neonatal complications.

Recommendations. The fetal heart rate should be monitored by a qualified individual before and after administration of neuraxial analgesia for labor. The Task Force recognizes that *continuous* electronic recording of the fetal heart rate may not be necessary in every clinical setting and may not be possible during initiation of neuraxial anesthesia.

II. Aspiration Prevention

Clear Liquids. There is insufficient published evidence to draw conclusions about the relationship between fasting times for clear liquids and the risk of emesis/reflux or pulmonary aspiration during labor. The consultants and ASA members both agree that oral intake of clear liquids during labor improves maternal comfort and satisfaction. Although the ASA members are equivocal, the consultants agree that oral intake of clear liquids during labor *does not* increase maternal complications.

Recommendations. The oral intake of modest amounts of clear liquids may be allowed for uncomplicated laboring patients. The uncomplicated patient undergoing elective cesarean delivery may have modest amounts of clear liquids up to 2 hours prior to induction of anesthesia. Examples of clear liquids include, but are not limited to, water, fruit juices without pulp, carbonated beverages, clear tea, black coffee, and sports drinks.** The volume of liquid ingested is less important than the presence of particulate matter in the liquid ingested. However, patients with additional risk factors for aspiration (*e.g.*, morbid obesity, diabetes, difficult airway), or patients at increased risk for operative delivery (*e.g.*, nonreassuring fetal heart rate pattern) may have further restrictions of oral intake, determined on a case-by-case basis.

** American Society of Anesthesiologists Task Force on Preoperative Fasting: Practice guidelines for preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration. *Anesthesiology* 1999; 90:896-905.

Solids. A specific fasting time for solids that is predictive of maternal anesthetic complications has not been determined. There is insufficient published evidence to address the safety of *any* particular fasting period for solids in obstetric patients. The consultants and ASA members both agree that the oral intake of solids during labor increases maternal complications. They both strongly agree that patients undergoing either elective cesarean delivery or postpartum tubal ligation should undergo a fasting period of 6 to 8 hours depending on the type of food ingested (*e.g.*, fat content).⁴ The Task Force recognizes that in laboring patients the timing of delivery is uncertain; therefore compliance with a predetermined fasting period before non-elective surgical procedures is not always possible.

Recommendations. Solid foods should be avoided in laboring patients. The patient undergoing elective surgery (*e.g.*, scheduled cesarean delivery or postpartum tubal ligation) should undergo a fasting period for solids of 6 to 8 hours depending on the type of food ingested (*e.g.*, fat content).⁴

Antacids, H₂ Receptor Antagonists, and Metoclopramide. The literature does not sufficiently examine the relationship between reduced gastric acidity and the frequency of emesis, pulmonary aspiration, morbidity, or mortality in obstetric patients who have aspirated gastric contents. Published evidence supports the efficacy of preoperative non-particulate antacids (*e.g.*, sodium citrate, sodium bicarbonate) in decreasing gastric acidity during the peripartum period. However, the literature is insufficient to examine the impact of non-particulate antacids on gastric volume. The literature suggests that H₂ receptor antagonists are effective in decreasing gastric acidity in obstetric patients, and supports the efficacy of metoclopramide in reducing peripartum nausea and vomiting. The consultants and ASA members agree that the administration of a non-particulate antacid prior to operative procedures reduces maternal complications.

Recommendations. Before surgical procedures (*i.e.*, cesarean delivery, postpartum tubal ligation), practitioners should consider the timely administration of non-particulate antacids, H₂ receptor antagonists, and/or metoclopramide for aspiration prophylaxis.

III. Anesthetic Care for Labor and Vaginal Delivery

Overview. Not all women require anesthetic care during labor or delivery. For women who request pain relief for labor and/or delivery, there are many effective analgesic techniques available. Maternal request represents sufficient justification for pain relief. In addition, maternal medical and obstetric conditions may warrant the provision of neuraxial techniques to improve maternal and neonatal outcome.

The choice of analgesic technique depends on the medical status of the patient, progress of labor, and resources at the facility. When sufficient resources (*e.g.*, anesthesia and nursing staff) are available, neuraxial catheter techniques should be one of the analgesic options offered. The choice of a specific neuraxial block should be individualized and based on anesthetic risk factors, obstetric risk factors, patient preferences, progress of labor, and resources at the facility.

When neuraxial catheter techniques are used for analgesia during labor or vaginal delivery, the primary goal is to provide adequate maternal analgesia with minimal motor block (*e.g.*, achieved with the administration of local anesthetics at low concentrations with or without opioids).

When a neuraxial technique is chosen, appropriate resources for the treatment of complications (*e.g.*, hypotension, systemic toxicity, high spinal anesthesia) should be available. If an opioid is added, treatments for related complications (*e.g.*, pruritus, nausea, respiratory depression) should be available. An intravenous infusion should be established before the initiation of neuraxial analgesia or anesthesia and maintained throughout the duration of the neuraxial analgesic or anesthetic. However, administration of a fixed volume of intravenous fluid is not required before neuraxial analgesia is initiated.

Timing of Neuraxial Analgesia and Outcome of Labor. Meta-analysis of the literature determined that the timing of neuraxial analgesia does not affect the frequency of cesarean delivery. The literature also suggests that other delivery outcomes (*i.e.*, spontaneous or instrumented) are also

unaffected. The consultants strongly agree and the ASA members agree that early initiation of epidural analgesia (*i.e.*, at cervical dilations of less than 5 cm vs equal to or greater than 5 cm) improves analgesia. They both *disagree* that motor block or maternal, fetal or neonatal side effects are increased by early administration.

Recommendations. Patients in early labor (*i.e.*, < 5 cm dilation) should be given the option of neuraxial analgesia when this service is available. Neuraxial analgesia should not be withheld on the basis of achieving an arbitrary cervical dilation, and should be offered on an individualized basis. Patients may be reassured that the use of neuraxial analgesia does not increase the incidence of cesarean delivery.

Neuraxial Analgesia and Trial of Labor After Prior Cesarean Delivery. Nonrandomized comparative studies suggest that epidural analgesia may be used in a trial of labor for previous cesarean delivery patients without adversely affecting the incidence of vaginal delivery. Randomized comparisons of epidural versus other anesthetic techniques were not found. The consultants and ASA members agree that neuraxial techniques improve the likelihood of vaginal delivery for patients attempting vaginal birth after cesarean delivery.

Recommendations: Neuraxial techniques should be offered to patients attempting vaginal birth after prior cesarean delivery. For these patients, it is also appropriate to consider early placement of a neuraxial catheter that can be used later for labor analgesia, or for anesthesia in the event of operative delivery.

Early Insertion of a Spinal or Epidural Catheter for Complicated Parturients. When caring for the complicated parturient, the literature is insufficient to assess whether the early insertion of a spinal or epidural catheter, with later administration of analgesia, improves maternal or neonatal outcomes. The consultants and ASA members agree that early insertion of a spinal or epidural catheter for complicated parturients reduces maternal complications.

Recommendations: Early insertion of a spinal or epidural catheter for obstetric (*e.g.*, twin gestation or preeclampsia) or anesthetic indications (*e.g.*, anticipated difficult airway or obesity) should be considered to reduce the need for general anesthesia if an emergent procedure becomes necessary. In these cases, the insertion of a spinal or epidural catheter may precede the onset of labor or a patient's request for labor analgesia.

Continuous Infusion Epidural Analgesia.

Continuous infusion epidural analgesia compared to parenteral opioids: The literature suggests that the use of continuous infusion epidural (CIE) local anesthetics with or without opioids provides greater quality of analgesia compared to parenteral (*i.e.*, intravenous or intramuscular) opioids. The consultants and ASA members strongly agree that CIE local anesthetics with or without opioids provide improved analgesia compared to parenteral opioids.

Meta-analysis of the literature indicates that there is a longer duration of labor, with an average duration of 24 min for the second stage, and a lower frequency of spontaneous vaginal delivery when continuous epidural local anesthetics are administered compared to *intravenous* opioids. Meta-analysis of the literature determined that there are no differences in the frequency of cesarean delivery. Neither the consultants nor ASA members agree that CIE local anesthetics compared to parenteral opioids significantly (1) increase the duration of labor, (2) decrease the chance of spontaneous delivery, (3) increase maternal side effects, or (4) increase fetal and neonatal side effects.

CIE compared to single-injection spinal: There is insufficient literature to assess the analgesic efficacy of CIE local anesthetics with or without opioids compared to *single-injection spinal opioids* with or without local anesthetics. The consultants are equivocal, but the ASA members agree that CIE local anesthetics improve analgesia compared to single-injection spinal opioids; both the consultants and ASA members are equivocal regarding the frequency of motor block. The consultants are equivocal, but the ASA members disagree that the use of CIE compared to single-injection spinal

opioids increases the duration of labor. They both *disagree* that CIE local anesthetics with or without opioids compared to single-injection spinal opioids with or without local anesthetics decreases the likelihood of spontaneous delivery or increases maternal, fetal or neonatal side effects.

CIE with and without opioids: The literature supports the *induction* of analgesia using epidural local anesthetics combined *with opioids* compared to equal concentrations of epidural local anesthetics *without opioids* for improved quality and longer duration of analgesia. The consultants strongly agree and the ASA members agree that the addition of opioids to epidural local anesthetics improves analgesia; they both disagree that fetal or neonatal side effects are increased. The consultants disagree, but the ASA members are equivocal regarding whether the addition of opioids increases maternal side effects.

The literature is insufficient to determine whether induction of analgesia using local anesthetics with opioids compared to *higher concentrations* of epidural local anesthetics without opioids provides improved quality or duration of analgesia. The consultants and ASA members are equivocal regarding improved analgesia, and they both disagree that maternal, fetal or neonatal side effects are increased using lower concentrations of epidural local anesthetics with opioids.

For *maintenance of analgesia*, the literature suggests that there are no differences in the analgesic efficacy of *low concentrations* of epidural local anesthetics with opioids compared to *higher concentrations* of epidural local anesthetics without opioids. The Task Force notes that the addition of an opioid to a local anesthetic infusion allows an even lower concentration of local anesthetic for providing equally effective analgesia. However, the literature is insufficient to examine whether a bupivacaine infusion concentration of *less than or equal to* 0.125% with an opioid provides comparable or improved analgesia compared to a bupivacaine concentration *greater than* 0.125%

without an opioid.^{††} Meta-analysis of the literature determined that low concentrations of epidural local anesthetics with opioids compared to higher concentrations of epidural local anesthetics without opioids are associated with reduced motor block. No differences in the duration of labor, mode of delivery, or neonatal outcomes are found when epidural local anesthetics with opioids are compared to epidural local anesthetics without opioids. The literature is insufficient to determine the effects of epidural local anesthetics with opioids on other maternal outcomes (*e.g.*, hypotension, nausea, pruritus, respiratory depression, urinary retention).

The consultants and ASA members both agree that maintenance of epidural analgesia using *low* concentrations of local anesthetics with opioids provides improved analgesia compared to *higher* concentrations of local anesthetics without opioids. The consultants agree, but the ASA members are equivocal regarding the improved likelihood of spontaneous delivery when lower concentrations of local anesthetics with opioids are used. The consultants strongly agree and the ASA members agree that motor block is reduced. They agree that maternal side effects are reduced with this drug combination. They are both equivocal regarding a reduction in fetal and neonatal side effects.

Recommendations. The selected analgesic/anesthetic technique should reflect patient needs and preferences, practitioner preferences or skills, and available resources. The continuous epidural infusion technique may be used for effective analgesia for labor and delivery. When a continuous epidural infusion of local anesthetic is selected, an opioid may be added to reduce the concentration of local anesthetic, improve the quality of analgesia, and minimize motor block.

Adequate analgesia for uncomplicated labor and delivery should be administered with the secondary goal of producing as little motor block as possible by using dilute concentrations of local anesthetics with opioids. The lowest concentration of local anesthetic infusion that provides adequate

^{††} References to bupivacaine are included for illustrative purposes only, and because bupivacaine is the most extensively studied local anesthetic for continuous infusion epidural analgesia. The Task Force recognizes that other local anesthetics are appropriate for continuous infusion epidural analgesia.

maternal analgesia and satisfaction should be administered. For example, an infusion concentration greater than 0.125% bupivacaine is unnecessary for labor analgesia in most patients.

Single-Injection Spinal Opioids With or Without Local Anesthetics. The literature suggests that spinal opioids with or without local anesthetics provide effective analgesia during labor without altering the incidence of neonatal complications. There is insufficient literature to compare spinal opioids with parenteral opioids. There is also insufficient literature to compare single-injection spinal opioids *with* local anesthetics *versus* single-injection spinal opioids *without* local anesthetics.

The consultants strongly agree and the ASA members agree that spinal opioids provide improved analgesia compared to parenteral opioids. They both disagree that, compared to parenteral opioids, spinal opioids increase the duration of labor, decrease the chance of spontaneous delivery, or increase fetal and neonatal side effects. The consultants are equivocal, but the ASA members disagree that maternal side effects are increased with spinal opioids.

Compared to spinal opioids *without* local anesthetics, the consultants and ASA members both agree that spinal opioids *with* local anesthetics provide improved analgesia. They both disagree that the chance of spontaneous delivery is decreased, and that fetal and neonatal side effects are increased. They are both equivocal regarding an increase in maternal side effects. However, they both agree that motor block is increased when local anesthetics are added to spinal opioids. Finally, the consultants disagree, but the ASA members are equivocal regarding an increase in the duration of labor.

Recommendations. Single-injection spinal opioids with or without local anesthetics may be used to provide effective, although time-limited, analgesia for labor when spontaneous vaginal delivery is anticipated. If labor is expected to last longer than the analgesic effects of the spinal drugs chosen, or if there is a good possibility of operative delivery, then a catheter technique instead of a single injection technique should be considered. A local anesthetic may be added to a spinal opioid to increase duration and improve quality of analgesia. The Task Force notes that the rapid onset of

analgesia provided by single-injection spinal techniques may be advantageous for selected patients (e.g., those in advanced labor).

Pencil-Point Spinal Needles. The literature supports the use of pencil-point spinal needles compared to cutting-bevel spinal needles to reduce the frequency of post-dural puncture headache. The consultants and ASA members both strongly agree that the use of pencil-point spinal needles reduces maternal complications.

Recommendations. Pencil-point spinal needles should be used instead of cutting-bevel spinal needles to minimize the risk of post-dural puncture headache.

Combined Spinal-Epidural Analgesia. The literature supports a faster onset time and equivalent analgesia with combined spinal-epidural (CSE) local anesthetics with opioids versus epidural local anesthetics with opioids. The literature is equivocal regarding the impact of CSE versus epidural local anesthetics with opioids on maternal satisfaction with analgesia, mode of delivery, hypotension, motor block, nausea, fetal heart rate changes, and Apgar scores. Meta-analysis of the literature indicates that the frequency of pruritus is increased with CSE.

The consultants and ASA members both agree that CSE local anesthetics with opioids provide improved early analgesia compared to epidural local anesthetics with opioids. They are equivocal regarding the impact of CSE with opioids on overall analgesic efficacy, duration of labor, and motor block. The consultants and ASA members both disagree that CSE increases the risk of fetal or neonatal side effects. The consultants disagree, but the ASA members are equivocal regarding whether CSE increases the incidence of maternal side effects.

Recommendations. CSE techniques may be used to provide effective and rapid onset of analgesia for labor.

Patient-Controlled Epidural Analgesia. The literature supports the efficacy of patient-controlled epidural analgesia (PCEA) *versus* CIE in providing equivalent analgesia with reduced drug

consumption. Meta-analysis of the literature indicates that the duration of labor is longer with PCEA compared to CIE for the first stage (*e.g.*, an average of 36 min) but not the second stage of labor. Meta-analysis of the literature also determined that mode of delivery, frequency of motor block, and Apgar scores are equivalent when PCEA administration is compared to CIE. The literature supports greater analgesic efficacy for PCEA with a background infusion compared to PCEA without a background infusion; meta-analysis of the literature also indicates no differences in the mode of delivery or frequency of motor block. The consultants and ASA members agree that PCEA compared to CIE improves analgesia and reduces the need for anesthetic interventions; they also agree that PCEA improves maternal satisfaction. The consultants and ASA members are equivocal regarding a reduction in motor block, an increased likelihood of spontaneous delivery, or a decrease in maternal side effects with PCEA compared to CIE. They both agree that PCEA with a background infusion improves analgesia, improves maternal satisfaction, and reduces the need for anesthetic intervention. The ASA members are equivocal, but the consultants disagree that a background infusion decreases the chance of spontaneous delivery or increases maternal side effects. The consultants and ASA members are equivocal regarding the effect of a background infusion on the incidence of motor block.

Recommendations. PCEA may be used to provide an effective and flexible approach for the maintenance of labor analgesia. The Task Force notes that the use of PCEA may be preferable to CIE for providing fewer anesthetic interventions, reduced dosages of local anesthetics, and less motor blockade than fixed-rate continuous epidural infusions. PCEA may be used with or without a background infusion.

IV. Removal of Retained Placenta

Anesthetic Techniques. The literature is insufficient to assess whether a particular type of anesthetic is more effective than another for removal of retained placenta. The consultants strongly agree and the ASA members agree that, if a functioning epidural catheter is in place, and the patient is

hemodynamically stable, epidural anesthesia is the preferred technique for the removal of retained placenta. The consultants and ASA members both agree that, in cases involving major maternal hemorrhage, general anesthesia is preferred over neuraxial anesthesia.

Recommendations. The Task Force notes that, in general, there is no preferred anesthetic technique for removal of retained placenta. However, if an epidural catheter is in place and the patient is hemodynamically stable, epidural anesthesia is preferable. Hemodynamic status should be assessed before administering neuraxial anesthesia. Aspiration prophylaxis should be considered.

Sedation/analgesia should be titrated carefully due to the potential risks of respiratory depression and pulmonary aspiration during the immediate postpartum period. In cases involving major maternal hemorrhage, general anesthesia with an endotracheal tube may be preferable to neuraxial anesthesia.

Uterine Relaxation. The literature suggests that nitroglycerin is effective for uterine relaxation during the removal of retained placenta. The consultants and ASA members both agree that the administration of nitroglycerin for uterine relaxation improves success in removing a retained placenta.

Recommendations. Nitroglycerin may be used as an alternative to terbutaline sulfate or general endotracheal anesthesia with halogenated agents for uterine relaxation during removal of retained placental tissue. Initiating treatment with incremental doses of intravenous or sublingual (*i.e.*, metered dose spray) nitroglycerin may relax the uterus sufficiently while minimizing potential complications (*e.g.*, hypotension).

V. Anesthetic Choices for Cesarean Delivery

Equipment, Facilities and Support Personnel. The literature is insufficient to evaluate the benefit of providing equipment, facilities and support personnel in the labor and delivery operating suite comparable to that available in the main operating suite. The consultants and ASA members strongly agree that the available equipment, facilities, and support personnel should be comparable.

Recommendations. Equipment, facilities, and support personnel available in the labor and delivery operating suite should be comparable to those available in the main operating suite. Resources for the treatment of potential complications (*e.g.*, failed intubation, inadequate analgesia, hypotension, respiratory depression, pruritus, vomiting) should also be available in the labor and delivery operating suite. Appropriate equipment and personnel should be available to care for obstetric patients recovering from major neuraxial or general anesthesia.

General, Epidural, Spinal, or Combined Spinal-Epidural Anesthesia. The literature suggests that induction-to-delivery times for general anesthesia (GA) are lower compared to epidural or spinal anesthesia, and that a higher frequency of maternal hypotension may be associated with epidural or spinal techniques. Meta-analysis of the literature found that Apgar scores at one and five minutes are lower for GA compared to epidural anesthesia, and suggests that Apgar scores are lower for GA *versus* spinal anesthesia. The literature is equivocal regarding differences in umbilical artery pH values when GA is compared with epidural or spinal anesthesia.

The consultants and ASA members agree that GA reduces the time to skin incision when compared to either epidural or spinal anesthesia; they also agree that GA increases maternal complications. The consultants are equivocal and the ASA members agree that GA increases fetal and neonatal complications. The consultants and ASA members both agree that epidural anesthesia increases the time to skin incision and decreases the quality of anesthesia compared to spinal anesthesia. They both disagree that epidural anesthesia increases maternal complications.

When spinal anesthesia is compared to epidural anesthesia, meta-analysis of the literature found that induction-to-delivery times are shorter for spinal anesthesia. The literature is equivocal regarding hypotension, umbilical pH values and Apgar scores. The consultants and ASA members agree that epidural anesthesia increases time to skin incision and reduces the quality of anesthesia when

compared to spinal anesthesia. They both disagree that epidural anesthesia increases maternal complications.

When CSE is compared to epidural anesthesia, meta-analysis of the literature found no differences in the frequency of hypotension or in 1 minute Apgar scores; the literature is insufficient to evaluate outcomes associated with the use of CSE compared to spinal anesthesia. The consultants and ASA members agree that CSE anesthesia improves anesthesia and reduces time to skin incision when compared to *epidural* anesthesia. The ASA members are equivocal, but the consultants disagree that maternal side effects are reduced. The consultants and ASA members both disagree that CSE improves anesthesia compared to *spinal* anesthesia. The ASA members are equivocal, but the consultants disagree that maternal side effects are reduced. The consultants strongly agree and the ASA members agree that CSE compared to spinal anesthesia increases flexibility of prolonged procedures, and they both agree that the time to skin incision is increased.

Recommendations. The decision to use a particular anesthetic technique for cesarean delivery should be individualized, based on several factors. These include anesthetic, obstetric or fetal risk factors (*e.g.*, elective versus emergency), the preferences of the patient, and the judgment of the anesthesiologist. Neuraxial techniques are preferred to general anesthesia for most cesarean deliveries. An indwelling epidural catheter may provide equivalent onset of anesthesia compared to initiation of spinal anesthesia for urgent cesarean delivery. If spinal anesthesia is chosen, pencil-point spinal needles should be used instead of cutting-bevel spinal needles. However, GA may be the most appropriate choice in some circumstances (*e.g.*, profound fetal bradycardia, ruptured uterus, severe hemorrhage, severe placental abruption). Uterine displacement (usually left displacement) should be maintained until delivery regardless of the anesthetic technique used.

Intravenous Fluid Preloading. The literature supports, and the consultants and ASA members agree that intravenous fluid preloading for spinal anesthesia reduces the frequency of maternal hypotension when compared to no fluid preloading.

Recommendations. Intravenous fluid preloading may be used to reduce the frequency of maternal hypotension following spinal anesthesia for cesarean delivery. Although fluid preloading reduces the frequency of maternal hypotension, initiation of spinal anesthesia should not be delayed in order to administer a fixed volume of intravenous fluid.

Ephedrine or Phenylephrine. The literature supports the administration of ephedrine and suggests that phenylephrine is effective in reducing maternal hypotension during neuraxial anesthesia for cesarean delivery. The literature is equivocal regarding the relative frequency of patients with breakthrough hypotension when infusions of ephedrine are compared to phenylephrine; however, lower umbilical cord pH values are reported after ephedrine administration. The consultants agree and the ASA members strongly agree that ephedrine is acceptable for treating hypotension during neuraxial anesthesia. The consultants strongly agree and the ASA members agree that phenylephrine is an acceptable agent for the treatment of hypotension.

Recommendations. Intravenous ephedrine and phenylephrine are both acceptable drugs for treating hypotension during neuraxial anesthesia. In the absence of maternal bradycardia, phenylephrine may be preferable because of improved fetal acid-base status in uncomplicated pregnancies.

Neuraxial Opioids for Postoperative Analgesia. For improved postoperative analgesia following cesarean delivery under epidural anesthesia, the literature supports the use of epidural opioids compared to intermittent injections of intravenous or intramuscular opioids. However, a higher frequency of pruritus was found with epidural opioids. The literature is insufficient to evaluate the impact of epidural opioids compared to intravenous PCA. In addition, the literature is insufficient to

evaluate spinal opioids compared to parenteral opioids. The consultants strongly agree and the ASA members agree that neuraxial opioids for postoperative analgesia improve analgesia and maternal satisfaction.

Recommendations. For postoperative analgesia after neuraxial anesthesia for cesarean delivery, neuraxial opioids are preferred over intermittent injections of parenteral opioids.

VI. Postpartum Tubal Ligation

There is insufficient literature to evaluate the benefits of neuraxial anesthesia compared to GA for postpartum tubal ligation. In addition, the literature is insufficient to evaluate the impact of the timing of a postpartum tubal ligation on maternal outcome. The consultants and ASA members both agree that neuraxial anesthesia for postpartum tubal ligation reduces complications compared to GA. The ASA members are equivocal but the consultants agree that a postpartum tubal ligation within 8 hours of delivery *does not* increase maternal complications.

Recommendations. For postpartum tubal ligation, the patient should have no oral intake of solid foods within 6 to 8 hours of the surgery, depending on the type of food ingested (*e.g.*, fat content).⁴ Aspiration prophylaxis should be considered. Both the timing of the procedure and the decision to use a particular anesthetic technique (*i.e.*, neuraxial versus general) should be individualized, based on anesthetic risk factors, obstetric risk factors (*e.g.*, blood loss), and patient preferences. However, neuraxial techniques are preferred to general anesthesia for most postpartum tubal ligations. The anesthesiologist should be aware that gastric emptying will be delayed in patients who have received opioids during labor, and that an epidural catheter placed for labor may be more likely to fail with longer post-delivery time intervals. If a postpartum tubal ligation is to be performed before the patient is discharged from the hospital, the procedure should not be attempted at a time when it might compromise other aspects of patient care on the labor and delivery unit.

VII. Management of Obstetric and Anesthetic Emergencies

Resources for Management of Hemorrhagic Emergencies. Descriptive studies and case reports suggest that the availability of resources for hemorrhagic emergencies may be associated with reduced maternal complications. The consultants and ASA members both strongly agree that the availability of resources for managing hemorrhagic emergencies reduces maternal complications.

Recommendations. Institutions providing obstetric care should have resources available to manage hemorrhagic emergencies (table 1). In an emergency, the use of type-specific or O negative blood is acceptable. In cases of intractable hemorrhage when banked blood is not available or the patient refuses banked blood, intraoperative cell-salvage should be considered if available.

Central Invasive Hemodynamic Monitoring. There is insufficient literature to examine whether pulmonary artery catheterization is associated with improved maternal, fetal or neonatal outcomes in patients with pregnancy-related hypertensive disorders. The literature is silent regarding the management of obstetric patients with central venous catheterization alone. The consultants and ASA members agree that the routine use of central venous or pulmonary artery catheterization does not reduce maternal complications in severe preeclamptic patients.

Recommendations. The decision to perform invasive hemodynamic monitoring should be individualized and based on clinical indications that include the patient's medical history and cardiovascular risk factors. The Task Force recognizes that not all practitioners have access to resources for utilization of central venous or pulmonary artery catheters in obstetric units.

Equipment for Management of Airway Emergencies. Case reports suggest that the availability of equipment for the management of airway emergencies may be associated with reduced maternal, fetal and neonatal complications. The consultants and ASA members both strongly agree that the immediate availability of equipment for the management of airway emergencies reduces maternal, fetal and neonatal complications.

Recommendations. Labor and delivery units should have personnel and equipment readily available to manage airway emergencies, to include a pulse oximeter and qualitative carbon dioxide detector, consistent with the ASA Practice Guidelines for Management of the Difficult Airway.^{‡‡} Basic airway management equipment should be immediately available during the provision of neuraxial analgesia (table 2). In addition, portable equipment for difficult airway management should be readily available in the operative area of labor and delivery units (table 3). The anesthesiologist should have a preformulated strategy for intubation of the difficult airway. When tracheal intubation has failed, ventilation with mask and cricoid pressure, or with a laryngeal mask airway or supraglottic airway device (*e.g.*, Combitube®, Intubating LMA [Fasttrach®]) should be considered for maintaining an airway and ventilating the lungs. If it is not possible to ventilate or awaken the patient, an airway should be created surgically.

Cardiopulmonary Resuscitation. The literature is insufficient to evaluate the efficacy of cardiopulmonary resuscitation in the obstetric patient during labor and delivery. In cases of cardiac arrest, the American Heart Association has stated that 4 to 5 minutes is the maximum time rescuers will have to determine if the arrest can be reversed by Basic Life Support and Advanced Cardiac Life Support interventions.^{§§} Delivery of the fetus may improve cardiopulmonary resuscitation of the mother by relieving aortocaval compression. The American Heart Association further notes that “the best survival rate for infants >24 to 25 weeks in gestation occurs when the delivery of the infant occurs no more than 5 minutes after the mother’s heart stops beating. This typically requires that the provider begin the hysterotomy about 4 minutes after cardiac arrest.”⁷ The consultants and ASA members both strongly agree that the immediate availability of basic and advanced life-support equipment in the labor and delivery suite reduces maternal, fetal and neonatal complications.

^{‡‡} American Society of Anesthesiologists Task Force on Management of the Difficult Airway: Practice guidelines for management of the difficult airway: An Updated Report. *Anesthesiology* 2003; 98:1269-1277.

^{§§} 2005 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2005; 112(22 and 24 Suppl):IV1-203.

Recommendations. Basic and advanced life-support equipment should be immediately available in the operative area of labor and delivery units. If cardiac arrest occurs during labor and delivery, standard resuscitative measures should be initiated. In addition, uterine displacement (usually left displacement) should be maintained. If maternal circulation is not restored within 4 min, cesarean delivery should be performed by the obstetrics team.

Appendix 1: Summary of Recommendations

I. Perianesthetic Evaluation

- Conduct a focused history and physical examination before providing anesthesia care
 - Maternal health and anesthetic history
 - Relevant obstetric history
 - Airway and heart and lung examination
 - Baseline blood pressure measurement
 - Back examination when neuraxial anesthesia is planned or placed
- A communication system should be in place to encourage early and ongoing contact between obstetric providers, anesthesiologists, and other members of the multidisciplinary team.
- Order or require a platelet count based on a patient's history, physical examination and clinical signs; a routine intrapartum platelet count is not necessary in the healthy parturient
- Order or require an intrapartum blood type and screen or cross-match based on maternal history, anticipated hemorrhagic complications (*e.g.*, placenta accreta in a patient with placenta previa and previous uterine surgery) and local institutional policies; a routine blood cross-match is not necessary for *healthy and uncomplicated* parturients
- The fetal heart rate should be monitored by a qualified individual before and after administration of neuraxial analgesia for labor; *continuous* electronic recording of the fetal heart rate may not be necessary in every clinical setting and may not be possible during initiation of neuraxial anesthesia

II. Aspiration Prophylaxis

- Oral intake of modest amounts of clear liquids may be allowed for uncomplicated laboring patients
- The uncomplicated patient undergoing elective cesarean delivery may have modest amounts of clear liquids up to 2 hours prior to induction of anesthesia
- The volume of liquid ingested is less important than the presence of particulate matter in the liquid ingested
- Patients with additional risk factors for aspiration (*e.g.*, morbid obesity, diabetes, difficult airway) or patients at increased risk for operative delivery (*e.g.*, nonreassuring fetal heart rate pattern) may have further restrictions of oral intake, determined on a case-by-case basis
- Solid foods should be avoided in laboring patients
- Patients undergoing elective surgery (*e.g.*, scheduled cesarean delivery or postpartum tubal ligation) should undergo a fasting period for solids of 6 to 8 hours depending on the type of food ingested (*e.g.*, fat content)
- Before surgical procedures (*i.e.*, cesarean delivery, postpartum tubal ligation) practitioners should consider timely administration of non-particulate antacids, H₂ receptor antagonists, and/or metoclopramide for aspiration prophylaxis

III. Anesthetic Care for Labor and Delivery

Neuraxial techniques-availability of resources

- When neuraxial techniques that include local anesthetics are chosen, appropriate resources for the treatment of complications (e.g., hypotension, systemic toxicity, high spinal anesthesia) should be available
- If an opioid is added, treatments for related complications (e.g., pruritus, nausea, respiratory depression) should be available
- An intravenous infusion should be established before the initiation of neuraxial analgesia or anesthesia and maintained throughout the duration of the neuraxial analgesic or anesthetic
- Administration of a fixed volume of intravenous fluid is not required before neuraxial analgesia is initiated

Timing of neuraxial analgesia and outcome of labor

- Neuraxial analgesia should not be withheld on the basis of achieving an arbitrary cervical dilation, and should be offered on an individualized basis when this service is available
- Patients may be reassured that the use of neuraxial analgesia does not increase the incidence of cesarean delivery

Neuraxial analgesia and trial of labor after prior cesarean delivery

- Neuraxial techniques should be offered to patients attempting vaginal birth after prior cesarean delivery
- For these patients, it is also appropriate to consider early placement of a neuraxial catheter that can be used later for labor analgesia or for anesthesia in the event of operative delivery

Early insertion of spinal or epidural catheter for complicated parturients

- Early insertion of a spinal or epidural catheter for obstetric (e.g., twin gestation or preeclampsia) or anesthetic indications (e.g., anticipated difficult airway or obesity) should be considered to reduce the need for general anesthesia if an emergent procedure becomes necessary
 - In these cases, the insertion of a spinal or epidural catheter may precede the onset of labor or a patient's request for labor analgesia

Continuous infusion epidural (CIE) analgesia

- The selected analgesic/anesthetic technique should reflect patient needs and preferences, practitioner preferences or skills, and available resources
- CIE may be used for effective analgesia for labor and delivery
- When a continuous epidural infusion of local anesthetic is selected, an opioid may be added to reduce the concentration of local anesthetic, improve the quality of analgesia, and minimize motor block
- Adequate analgesia for uncomplicated labor and delivery should be administered with the secondary goal of producing as little motor block as possible by using dilute concentrations of local anesthetics with opioids
- The lowest concentration of local anesthetic infusion that provides adequate maternal analgesia and satisfaction should be administered

Single-injection spinal opioids with or without local anesthetics

- Single-injection spinal opioids with or without local anesthetics may be used to provide effective, although time-limited, analgesia for labor when spontaneous vaginal delivery is anticipated
- If labor is expected to last longer than the analgesic effects of the spinal drugs chosen, or if there is a good possibility of operative delivery, then a catheter technique instead of a single injection technique should be considered
- A local anesthetic may be added to a spinal opioid to increase duration and improve quality of analgesia

Pencil-point spinal needles

- Pencil-point spinal needles should be used instead of cutting-bevel spinal needles to minimize the risk of post dural puncture headache

Combined spinal-epidural (CSE) anesthetics

- CSE techniques may be used to provide effective and rapid analgesia for labor

Patient-controlled epidural analgesia (PCEA)

- PCEA may be used to provide an effective and flexible approach for the maintenance of labor analgesia
- PCEA may be preferable to CIE for providing fewer anesthetic interventions, reduced dosages of local anesthetics, and less motor blockade than fixed-rate continuous epidural infusions.
- PCEA may be used with or without a background infusion

IV. Removal of Retained Placenta

- In general, there is no preferred anesthetic technique for removal of retained placenta
 - If an epidural catheter is in place and the patient is hemodynamically stable, epidural anesthesia is preferable
- Hemodynamic status should be assessed before administering neuraxial anesthesia
- Aspiration prophylaxis should be considered
- Sedation/analgesia should be titrated carefully due to the potential risks of respiratory depression and pulmonary aspiration during the immediate postpartum period
- In cases involving major maternal hemorrhage, general anesthesia with an endotracheal tube may be preferable to neuraxial anesthesia
- Nitroglycerin may be used as an alternative to terbutaline sulfate or general endotracheal anesthesia with halogenated agents for uterine relaxation during removal of retained placental tissue
 - Initiating treatment with incremental doses of intravenous or sublingual (i.e., metered dose spray) nitroglycerin may relax the uterus sufficiently while minimizing potential complications (e.g., hypotension)

V. Anesthetic Choices for Cesarean Delivery

- Equipment, facilities, and support personnel available in the labor and delivery operating suite should be comparable to those available in the main operating suite
 - Resources for the treatment of potential complications (*e.g.*, failed intubation, inadequate analgesia, hypotension, respiratory depression, pruritus, vomiting) should be available in the labor and delivery operating suite
 - Appropriate equipment and personnel should be available to care for obstetric patients recovering from major neuraxial or general anesthesia
- The decision to use a particular anesthetic technique should be individualized based on anesthetic, obstetric or fetal risk factors (*e.g.*, elective versus emergency), the preferences of the patient, and the judgment of the anesthesiologist
 - Neuraxial techniques are preferred to general anesthesia for most cesarean deliveries
- An indwelling epidural catheter may provide equivalent onset of anesthesia compared to initiation of spinal anesthesia for urgent cesarean delivery
- If spinal anesthesia is chosen, pencil-point spinal needles should be used instead of cutting-bevel spinal needles
- GA may be the most appropriate choice in some circumstances (*e.g.*, profound fetal bradycardia, ruptured uterus, severe hemorrhage, severe placental abruption)
- Uterine displacement (usually left displacement) should be maintained until delivery regardless of the anesthetic technique used
- Intravenous fluid preloading may be used to reduce the frequency of maternal hypotension following spinal anesthesia for cesarean delivery
- Initiation of spinal anesthesia should not be delayed in order to administer a fixed volume of intravenous fluid
- Intravenous ephedrine and phenylephrine are both acceptable drugs for treating hypotension during neuraxial anesthesia
 - In the absence of maternal bradycardia, phenylephrine may be preferable because of improved fetal acid-base status in uncomplicated pregnancies
- For postoperative analgesia after neuraxial anesthesia for cesarean delivery, neuraxial opioids are preferred over intermittent injections of parenteral opioids

VI. Postpartum Tubal Ligation

- For postpartum tubal ligation, the patient should have no oral intake of solid foods within 6 to 8 hours of the surgery, depending on the type of food ingested (*e.g.*, fat content)
- Aspiration prophylaxis should be considered
- Both the timing of the procedure and the decision to use a particular anesthetic technique (*i.e.*, neuraxial versus general) should be individualized, based on anesthetic risk factors, obstetric risk factors (*e.g.*, blood loss), and patient preferences
- Neuraxial techniques are preferred to general anesthesia for most postpartum tubal ligations
 - Be aware that gastric emptying will be delayed in patients who have received opioids during labor, and that an epidural catheter placed for labor may be more likely to fail with longer post-delivery time intervals
- If a postpartum tubal ligation is to be performed before the patient is discharged from the hospital, the procedure should not be attempted at a time when it might compromise other aspects of patient care on the labor and delivery unit

VII. Management of Obstetric and Anesthetic Emergencies

- Institutions providing obstetric care should have resources available to manage hemorrhagic emergencies
 - In an emergency, the use of type-specific or O negative blood is acceptable
 - In cases of intractable hemorrhage when banked blood is not available or the patient refuses banked blood, intraoperative cell-salvage should be considered if available
 - The decision to perform invasive hemodynamic monitoring should be individualized and based on clinical indications that include the patient's medical history and cardiovascular risk factors
- Labor and delivery units should have personnel and equipment readily available to manage airway emergencies, to include a pulse oximeter and qualitative carbon dioxide detector, consistent with the ASA Practice Guidelines for Management of the Difficult Airway
 - Basic airway management equipment should be immediately available during the provision of neuraxial analgesia
 - Portable equipment for difficult airway management should be readily available in the operative area of labor and delivery units
 - The anesthesiologist should have a preformulated strategy for intubation of the difficult airway
 - When tracheal intubation has failed, ventilation with mask and cricoid pressure, or with a laryngeal mask airway or supraglottic airway device (*e.g.*, Combitube®, Intubating LMA [Fastrach®]) should be considered for maintaining an airway and ventilating the lungs
 - If it is not possible to ventilate or awaken the patient, an airway should be created surgically
- Basic and advanced life-support equipment should be immediately available in the operative area of labor and delivery units
- If cardiac arrest occurs during labor and delivery, standard resuscitative measures should be initiated
 - Uterine displacement (usually left displacement) should be maintained
 - If maternal circulation is not restored within 4 min, cesarean delivery should be performed by the obstetrics team

Table 1. Suggested Resources for Obstetric Hemorrhagic Emergencies*

- Large bore intravenous catheters
- Fluid warmer
- Forced air body warmer
- Availability of blood bank resources
- Equipment for infusing intravenous fluids and blood products rapidly. Examples include, but are not limited to, hand squeezed fluid chambers, hand inflated pressure bags, and automatic infusion devices.

* The items listed represent suggestions. The items should be customized to meet the specific needs, preferences, and skills of the practitioner and health-care facility.

Table 2. Suggested Resources for Airway Management During Initial Provision of Neuraxial Anesthesia*

- Laryngoscope and assorted blades
- Endotracheal tubes, with stylets
- Oxygen source
- Suction source with tubing and catheters
- Self-inflating bag and mask for positive pressure ventilation.
- Medications for blood pressure support, muscle relaxation, and hypnosis
- Qualitative carbon dioxide detector
- Pulse oximeter

* The items listed represent suggestions. The items should be customized to meet the specific needs, preferences, and skills of the practitioner and health-care facility.

Table 3. Suggested Contents of a Portable Storage Unit for Difficult Airway Management for Cesarean Section Rooms¹

- Rigid laryngoscope blades of alternate design and size from those routinely used
- Laryngeal mask airway
- Endotracheal tubes of assorted size
- Endotracheal tube guides. Examples include (but are not limited to) semi-rigid stylets with or without a hollow core for jet ventilation, light wands, and forceps designed to manipulate the distal portion of the endotracheal tube
- Retrograde intubation equipment
- At least one device suitable for emergency non-surgical airway ventilation. Examples include (but are not limited to) a hollow jet ventilation stylet with a transtracheal jet ventilator, and a supraglottic airway device (e.g., Combitube[®], Intubating LMA [Fastrach[®]])
- Fiberoptic intubation equipment
- Equipment suitable for emergency surgical airway access (e.g., cricothyrotomy)
- An exhaled carbon dioxide detector
- Topical anesthetics and vasoconstrictors

¹ Adapted from the Practice Guidelines for Management of the Difficult Airway: An Updated Report by the American Society of Anesthesiologists. *Anesthesiology* 2003; 98:1269-1277. The items listed represent suggestions. The items should be customized to meet the specific needs, preferences, and skills of the practitioner and health-care facility.

Appendix 2: Methods and Analyses

The scientific assessment of these Guidelines was based on evidence linkages or statements regarding potential relationships between clinical interventions and outcomes. The interventions listed below were examined to assess their impact on a variety of outcomes related to obstetric anesthesia.^{***}

1. Perianesthetic Evaluation

- i.* A directed history and physical examination
- ii.* Communication between anesthetic and obstetric providers
- iii.* A routine intrapartum platelet count does not reduce maternal anesthetic complications.
- iv.* For suspected preeclampsia or coagulopathy an intrapartum platelet count
- v.* An intrapartum blood type & screen for all parturients reduces maternal complications
- vi.* For healthy and uncomplicated parturients, a blood cross-match is unnecessary
- vii.* Perianesthetic recording of the fetal heart rate reduces fetal and neonatal complications

2. Aspiration Prophylaxis in the Obstetric Patient

- i.* Oral intake of clear liquids during labor improves patient comfort and satisfaction, but does not increase maternal complications
- ii.* Oral intake of solids during labor increases maternal complications
- iii.* A fasting period for solids of 6 to 8 hours before an elective cesarean reduces maternal complications
- iv.* Non-particulate antacids versus no antacids prior to operative procedures (excluding operative vaginal delivery) reduces maternal complications.

3. Anesthetic Care for Labor and Delivery^{†††}

- i.* Neuraxial Techniques
 - a.* Prophylactic spinal or epidural catheter insertion for complicated parturients reduces maternal complications
 - b.* Continuous epidural infusion of local anesthetics with or without opioids versus parenteral opioids
 - c.* Continuous epidural infusion of local anesthetics with or without opioids versus spinal opioids with or without local anesthetics
 - d.* Induction of epidural analgesia using local anesthetics with opioids versus equal concentrations of epidural local anesthetics without opioids
 - e.* Induction of epidural analgesia using local anesthetics with opioids versus higher concentrations of epidural local anesthetics without opioids
 - f.* Maintenance of epidural infusion of lower concentrations of local anesthetics with opioids versus higher concentrations of local anesthetics without opioids (e.g., bupivacaine concentrations < 0.125% with opioids versus concentrations > 0.125% without opioids)

^{***} Unless otherwise specified, outcomes for the listed interventions refer to the reduction of maternal, fetal and neonatal complications.

^{†††} Additional outcomes include improved analgesia, analgesic use, maternal comfort and satisfaction

- g. Single-injection spinal opioids with or without local anesthetics versus parenteral opioids
 - h. Single-injection spinal opioids with local anesthetics versus spinal opioids without local anesthetics
- ii. Combined Spinal-Epidural (CSE) Techniques
 - a. CSE local anesthetics with opioids versus epidural local anesthetics with opioids.
 - iii. Patient-Controlled Epidural Analgesia (PCEA)
 - a. PCEA versus continuous infusion epidurals
 - b. PCEA with a background infusion versus PCEA without a background infusion
 - iv. Neuraxial Analgesia, Timing of Initiation, and Progress of Labor
 - a. Administering epidural analgesia at cervical dilations of < 5 centimeters (versus > 5 cm)
 - b. Neuraxial techniques for patients attempting vaginal birth after prior cesarean delivery

4. Removal of Retained Placenta

- i. If an epidural catheter is *in situ* and the patient is hemodynamically stable, epidural anesthesia is preferred over general or spinal anesthesia to improve the success at removing retained placenta.
- ii. In cases involving major maternal hemorrhage, general anesthesia is preferred over neuraxial anesthesia to reduce maternal complications.
- iii. Administration of nitroglycerin for uterine relaxation improves success at removing retained placenta.

5. Anesthetic Choices for Cesarean Delivery

- i. Equipment, facilities, and support personnel available in the labor and delivery suite should be comparable to that available in the main operating suite
- ii. General anesthesia versus epidural anesthesia
- iii. General anesthesia versus spinal anesthesia
- iv. Epidural anesthesia versus spinal anesthesia
- v. CSE anesthesia versus epidural anesthesia
- vi. CSE anesthesia versus spinal anesthesia
- vii. Use of pencil-point spinal needles versus cutting-bevel spinal needles reduces maternal complications
- viii. Intravenous fluid preloading versus no intravenous fluid preloading for spinal anesthesia reduces maternal hypotension
- ix. Ephedrine or phenylephrine reduces maternal hypotension during neuraxial anesthesia
- x. Neuraxial opioids versus parenteral opioids for postoperative analgesia after neuraxial anesthesia for cesarean delivery

6. Postpartum Tubal Ligation

- i. Neuraxial anesthesia versus general anesthesia
- ii. A postpartum tubal ligation within 8 hours of delivery does not increase maternal complications

7. Management of Complications

- i. Availability of resources for management of hemorrhagic emergencies
- ii. Immediate availability of equipment for management of airway emergencies
- iii. Immediate availability of basic and advanced life-support equipment in the labor and delivery suite.
- iv. Invasive hemodynamic monitoring for severe preeclamptic patients.

Scientific evidence was derived from aggregated research literature, and opinion-based evidence was obtained from surveys, open presentations and other activities (*e.g.*, internet posting). For purposes of literature aggregation, potentially relevant clinical studies were identified *via* electronic and manual searches of the literature. The electronic and manual searches covered a 67-year period from 1940 through 2006. More than 4000 citations were initially identified, yielding a total of 2986 non-overlapping articles that addressed topics related to the evidence linkages. After review of the articles, 2549 studies did not provide direct evidence, and were subsequently eliminated. A total of 437 articles contained direct linkage-related evidence.

Initially, each pertinent outcome reported in a study was classified as supporting an evidence linkage, refuting a linkage, or equivocal. The results were then summarized to obtain a directional assessment for each evidence linkage before conducting a formal meta-analysis. Literature pertaining to 11 evidence linkages contained enough studies with well-defined experimental designs and statistical information sufficient for meta-analyses. These linkages were: (1) non-particulate antacids *versus* no antacids, (2) continuous epidural infusion of local anesthetics with or without opioids *versus* parenteral opioids, (3) induction of epidural analgesia using local anesthetics with opioids *versus* equal concentrations of epidural local anesthetics without opioids, (4) maintenance of epidural infusion of lower concentrations of local anesthetics with opioids *versus* higher concentrations of local anesthetics without opioids, (5) CSE local anesthetics with opioids *versus* epidural local anesthetics with opioids, (6) PCEA *versus* continuous infusion epidurals, (7) general anesthesia *versus* epidural anesthesia for cesarean delivery, (8), CSE anesthesia *versus* epidural anesthesia for cesarean delivery,

(9), use of pencil-point spinal needles *versus* cutting-bevel spinal needles, (10), ephedrine or phenylephrine reduces maternal hypotension during neuraxial anesthesia, and (11) neuraxial opioids *versus* parenteral opioids for postoperative analgesia after neuraxial anesthesia for cesarean delivery.

General variance-based effect-size estimates or combined probability tests were obtained for continuous outcome measures, and Mantel-Haenszel odds-ratios were obtained for dichotomous outcome measures. Two combined probability tests were employed as follows: (1) the Fisher Combined Test, producing chi-square values based on logarithmic transformations of the reported p-values from the independent studies, and (2) the Stouffer Combined Test, providing weighted representation of the studies by weighting each of the standard normal deviates by the size of the sample. An odds-ratio procedure based on the Mantel-Haenszel method for combining study results using 2 x 2 tables was used with outcome frequency information. An acceptable significance level was set at $p < 0.01$ (one-tailed). Tests for heterogeneity of the independent studies were conducted to assure consistency among the study results. DerSimonian-Laird random-effects odds ratios were obtained when significant heterogeneity was found ($p < 0.01$). To control for potential publishing bias, a "fail-safe n" value was calculated. No search for unpublished studies was conducted, and no reliability tests for locating research results were done.

Meta-analytic results are reported in table 4. To be accepted as significant findings, Mantel-Haenszel odds-ratios must agree with combined test results whenever both types of data are assessed. In the absence of Mantel-Haenszel odds-ratios, findings from both the Fisher and weighted Stouffer combined tests must agree with each other to be acceptable as significant.

Interobserver agreement among Task Force members and two methodologists was established by interrater reliability testing. Agreement levels using a kappa (k) statistic for two-rater agreement pairs were as follows: (1) type of study design, $k = 0.83-0.94$; (2) type of analysis, $k = 0.71-0.93$; (3) evidence linkage assignment, $k = 0.87-1.00$; and (4) literature inclusion for database, $k = 0.74-1.00$.

Three-rater chance-corrected agreement values were: (1) study design, $Sav = 0.884$, $Var(Sav) = 0.004$; (2) type of analysis, $Sav = 0.805$, $Var(Sav) = 0.009$; (3) linkage assignment, $Sav = 0.911$, $Var(Sav) = 0.002$; (4) literature database inclusion, $Sav = 0.660$, $Var(Sav) = 0.024$. These values represent moderate to high levels of agreement.

Consensus was obtained from multiple sources, including: (1) survey opinion from consultants who were selected based on their knowledge or expertise in obstetric anesthesia or maternal and fetal medicine, (2) survey opinions solicited from active members of the ASA, (3) testimony from attendees of publicly-held open forums at two national anesthesia meetings, (4) Internet commentary, and (5) Task Force opinion and interpretation. The survey rate of return was 75% ($n = 76$ of 102) for the consultants, and 2326 surveys were received from active ASA members. Results of the surveys are reported in tables 5 and 6, and in the text of the Guidelines.

The consultants were asked to indicate which, if any, of the evidence linkages would change their clinical practices if the Guidelines were instituted. The rate of return was 35% ($n = 36$). The percent of responding Consultants expecting *no change* associated with each linkage were as follows: perianesthetic evaluation - 97%; aspiration prophylaxis- 83%; anesthetic care for labor and delivery - 89%; removal of retained placenta - 97%; anesthetic choices for cesarean delivery - 97%; postpartum tubal ligation - 97%; and management of complications - 94%. Ninety-seven percent of the respondents indicated that the Guidelines would have *no effect* on the amount of time spent on a typical case. One respondent indicated that there would be an increase of 5 minutes in the amount of time spent on a typical case with the implementation of these Guidelines.

Table 4. Meta-Analysis Summary

Linkages	N	Fisher Chi-square	p	Weighted Stouffer Zc	p	Effect Size	Mantel-Haenszel OR	CI
ASPIRATION PROPHYLAXIS								
Non-particulate antacids vs no antacids								
Gastric pH ¹	5	66.80	0.001	9.78	0.001	0.88	-	-
Metoclopramide vs no metoclopramide								
Nausea	6	-	-	-	-	-	0.25	
		0.14-0.46	-	ns				
Vomiting	6	-	-	-	-	-	0.36	
		0.19-0.68	-	ns				
ANESTHETIC CARE FOR LABOR AND VAGINAL DELIVERY								
CIE local anesthetics +/- opioids vs IV opioids								
Duration of labor 1 st stage	5	50.19	0.001	5.42	0.001	0.15	-	-
Duration of labor 2 nd stage	7	67.53	0.001	4.84	0.001	0.21	-	-
Spontaneous delivery	8	-	-	-	-	-	0.53	
		0.42-0.68	-	ns				
Cesarean delivery ²	8	-	-	-	-	-	0.88	
		0.50-1.47	-	0.01				
Fetal acidosis	5	-	-	-	-	-	0.71	
		0.51-0.98	-	ns				
1 min Apgar	5	-	-	-	-	-	1.62	
		1.03-2.54	-	ns				
5 min Apgar	5	-	-	-	-	-	1.17	
		0.41-3.32	-	ns				
Epidural induction LA+O vs equal LA doses								
Analgesia (mean, SD)	6	91.21	0.001	17.70	0.001	0.99	-	-
Analgesia (pain relief)	5	-	-	-	-	-	4.03	
		2.14-7.56	-	ns				
Duration of labor	5	38.62	0.001	0.04	0.480	0.01	-	-
Spontaneous delivery	8	-	-	-	-	-	0.97	
		0.69-1.35	-	ns				
Hypotension	8	-	-	-	-	-	0.79	
		0.44-1.44	-	ns				
Motor block ¹	5	-	-	-	-	-	0.44	
		0.24-0.81	-	ns				
Nausea or vomiting	5	-	-	-	-	-	1.22	
		0.46-3.24	-	ns				
Pruritus	8	-	-	-	-	-	5.98	
		3.15-11.35	-	ns				
1 min Apgar	7	-	-	-	-	-	0.83	
		0.45-1.53	-	ns				
5 min Apgar	7	-	-	-	-	-	1.60	
		0.45-5.73	-	ns				

Duration of labor	5	19.82	0.030	1.99	0.020	0.05	-	-
Spontaneous delivery	8	-	-	-	-	-	1.08	-
		0.82-1.42	-	ns				
Motor block	6	-	-	-	-	-	0.29	-
		0.21-0.40	-	ns				
1 min Apgar	6	-	-	-	-	-	0.94	-
		0.60-1.47	-	ns				
5 min Apgar	6	-	-	-	-	-	2.62	-
		0.75-9.19	-	ns				
Pencil-point vs cutting-bevel spinal needles								
Post-dural puncture headache	5	-	-	-	-	-	0.34	-
		0.18-0.63	-	ns				
CSE LA+O vs epidural LA+O								
Analgesia (pain relief) ²	7	-	-	-	-	-	1.16	-
		0.62-1.85	-	0.010				
Satisfaction with analgesia	5	-	-	-	-	-	1.45	-
		0.89-2.34	-	ns				
Analgesia (time to onset)	5	57.80	0.001	-13.33	0.001	0.90	-	-
Spontaneous delivery	13	-	-	-	-	-	0.99	-
		0.85-1.15	-	ns				
Hypotension	8	-	-	-	-	-	1.71	-
		0.72-4.03	-	ns				
Motor block	7	-	-	-	-	-	1.20	-
		0.90-1.60	-	ns				
Nausea	5	-	-	-	-	-	1.22	-
		0.63-2.36	-	ns				
Pruritus ²	9	-	-	-	-	-	4.86	-
		1.63-14.65	-	0.001				
Motor block	7	-	-	-	-	-	1.20	-
		0.90-1.60	-	ns				
Fetal heart rate changes	6	-	-	-	-	-	1.25	-
		0.92-1.70	-	ns				
1 min Apgar	7	-	-	-	-	-	1.16	-
		0.76-1.77	-	ns				
5 min Apgar	8	-	-	-	-	-	1.33	-
		0.53-3.37	-	ns				
PCEA versus CIE								
Pain relief/score	5	21.78	0.020	0.17	0.433	0.04	-	-
Analgesic use	7	84.98	0.001	10.74	0.001	0.85	-	-
Duration of labor 1 st stage	5	42.42	0.001	5.24	0.001	0.44	-	-
Duration of labor 2 nd stage	6	43.08	0.001	2.01	0.022	0.18	-	-
Spontaneous delivery	13	-	-	-	-	-	1.22	-
		0.83-1.79	-	ns				
Motor block ²	7	-	-	-	-	-	0.52	-
		0.15-3.44	-	0.010				
1 min Apgar	6	-	-	-	-	-	0.63	-
		0.27-1.50	-	ns				

PCEA with background infusion versus PCEA

Analgesia (pain relief)	5	-	-	-	-	-	3.33	
	1.87-5.92			ns				
Spontaneous delivery	5	-	-	-	-	-	0.83	
	0.41-1.69			ns				
Motor block	5	-	-	-	-	-	1.18	
	0.47-2.97			ns				

Early versus late epidural

Cesarean delivery	5	-	-	-	-	-	0.95	
	0.67-1.35			ns				

ANESTHETIC CHOICES FOR CESAREAN DELIVERY**GA vs epidural**

Umbilical pH	5	49.04	0.001	0.52	0.300	0.37	-	-
1 min Apgar	5	49.04	0.001	-2.72	0.003	0.01	-	-
5 min Apgar	5	28.40	0.005	-2.95	0.002	0.08	-	-

CSE vs epidural

Hypotension	5	-	-	-	-	-	0.92	
	0.44-1.94			ns				
Umbilical pH	5	55.91	0.001	1.80	0.036	0.11	-	-
1 min Apgar	5	-	-	-	-	-	0.55	
	0.22-1.52			ns				

Fluid preloading vs no preloading

Hypotension ¹	6	-	-	-	-	-	0.46	
	0.29-0.73			ns				

Ephedrine vs placebo

Hypotension	7	-	-	-	-	-	0.26	
	0.14-0.48			ns				

Ephedrine vs phenylephrine

Hypotension	6	-	-	-	-	-	1.74	
	0.97-3.12			ns				
Umbilical pH	6	59.68	0.001	-7.55	0.001	0.71	-	-

Neuraxial vs parenteral O for postoperative analgesia

Analgesia	7	75.12	0.001	5.82	0.001	0.61	-	-
Nausea	9	-	-	-	-	-	1.13	
	0.57-2.22			ns				
Vomiting	6	-	-	-	-	-	1.02	
	0.37-2.81			ns				

¹ Non-randomized comparative studies included in analysis

² Dersimonian-Laird random effects odds ratio

OR = odds ratio; CIE = continuous infusion epidural; IV = intravenous; LA = local anesthetics; O = opioids;
LA+O = local anesthetics with opioids; SD = standard deviation; CSE = combined spinal epidural; PCEA =
patient-controlled epidural analgesia; GA = general anesthesia

Table 5. Consultant Survey Responses ^{†††}

	N	Percent Responding to Each Item				
		Strongly Agree	Agree	Uncertain	Disagree	Strongly Disagree
Perianesthetic Evaluation:						
1. Directed history and physical exam reduces maternal, fetal and neonatal complications	76	72.4*	26.3	1.3	0.0	0.0
2. Communication between anesthetic and obstetric providers reduces maternal, fetal and neonatal complications	76	89.5*	10.5	0.0	0.0	0.0
3. A routine intrapartum platelet count does not reduce maternal anesthetic complications	75	36.0	44.0*	8.0	10.7	1.3
4. An intrapartum platelet count reduces maternal anesthetic complications:						
For suspected preeclampsia	76	46.1	36.8*	9.2	7.9	0.0
For suspected coagulopathy	76	59.2*	32.9	5.3	2.6	0.0
5. All parturients should have an intrapartum blood sample sent to the blood bank to reduce maternal complications	76	21.1	32.9*	17.1	26.3	2.6
6. Perianesthetic recording of the fetal heart rate reduces fetal and neonatal complications	76	18.4	59.2*	13.2	9.2	0.0
Aspiration Prophylaxis:						
7a. Oral intake of clear liquids <i>during</i> labor improves patient comfort and satisfaction	76	32.9	60.5*	1.3	3.9	1.3
7b. Oral intake of clear liquids <i>during labor</i> does not increase maternal complications	75	16.0	45.3*	22.7	12.0	4.0
8a. Oral intake of solids <i>during labor</i> increases maternal complications	76	47.4	32.9*	10.5	5.3	3.9
8b. The patient undergoing elective cesarean delivery should undergo a fasting period for solids of 6 to 8 hours depending on the type of food ingested (e.g., fat content)	76	65.8*	30.3	3.9	0.0	0.0
8c. The patient undergoing elective postpartum tubal ligation should undergo a fasting period for solids of 6 to 8 hours depending on the type of food ingested (e.g., fat content)	76	56.6*	27.6	9.2	5.3	1.3
9. Administration of a nonparticulate antacid prior to operative procedures reduces maternal complications	75	29.3	45.3*	18.7	5.3	1.3
Anesthetic Care for Labor and Delivery:						
 Neuraxial Techniques:						
10. Prophylactic spinal or epidural catheter insertion for complicated parturients reduces maternal complications	75	42.7	40.0*	16.0	1.3	0.0

^{†††} N = the number of consultants who responded to each item. An asterisk beside a percentage score indicates the median.

	N	Percent Responding to Each Item				
		Strongly Agree	Agree	Uncertain	Disagree	Strongly Disagree
11. Continuous epidural infusion using local anesthetics with or without opioids versus parenteral opioids:						
Improves analgesia	75	84.0*	16.0	0.0	0.0	0.0
Increases the duration of labor	75	4.0	24.0	21.3	36.0*	14.7
Decreases the chance of spontaneous delivery	74	4.1	16.2	12.2	41.9*	25.7
Increases maternal side-effects	75	1.3	8.0	14.7	42.7*	33.3
Increases fetal and neonatal side-effects	75	0.0	4.0	6.7	46.7*	42.7
12. Continuous epidural infusion using local anesthetics with or without opioids versus spinal opioids with or without local anesthetics:						
Improves analgesia	74	12.2	25.7	20.3*	35.1	6.8
Increases the duration of labor	75	0.0	16.0	37.3*	34.7	12.0
Decreases the chance of spontaneous delivery	73	0.0	9.6	26.0	45.2*	19.2
Increases maternal motor block	74	5.4	41.9	17.6*	28.4	6.8
Increases maternal side-effects	74	0.0	6.8	27.0	52.7*	13.5
Increases fetal and neonatal side-effects	75	0.0	1.3	21.3	52.0*	25.3
13a. Induction of epidural analgesia using local anesthetics with opioids versus epidural analgesia with equal concentrations of local anesthetics without opioids:						
Improves analgesia	74	54.1*	39.2	1.4	4.1	1.4
Increases maternal side-effects	74	6.8	28.4	10.8	45.9*	8.1
Increases fetal and neonatal side-effects	74	0.0	2.7	12.2	59.5*	25.7
13b. Induction of epidural analgesia using low-dose local anesthetics with opioids versus higher concentrations of epidural local anesthetics without opioids:						
Improves analgesia	74	23.0	21.6	21.6*	32.4	1.4
Increases maternal side-effects	74	0.0	10.8	12.2	50.0*	27.0
Increases fetal and neonatal side-effects	74	0.0	2.7	17.6	52.7*	27.0
14a. Maintenance of epidural infusion of lower concentrations of local anesthetics with opioids versus higher concentrations of local anesthetics without opioids:						
Improves analgesia	74	21.6	28.4*	27.0	23.0	0.0
Reduces the duration of labor	74	4.1	35.1	40.5*	17.6	2.7
Improves the chance of spontaneous delivery	74	12.2	60.8*	14.9	10.8	1.4
Reduces maternal motor block	74	51.4*	43.2	5.4	0.0	0.0
Reduces maternal side-effects	74	16.2	44.6*	23.0	16.2	0.0
Reduces fetal and neonatal side-effects	74	8.1	24.3	32.4*	32.4	2.7

	N	Percent Responding to Each Item				
		Strongly Agree	Agree	Uncertain	Disagree	Strongly Disagree
14b. Maintenance of epidural analgesia using bupivacaine \leq 0.125% with opioids versus bupivacaine concentrations $>$ 0.125% without opioids:						
Improves analgesia	74	21.6	33.8*	21.6	23.0	0.0
Reduces the duration of labor	74	6.8	33.8	45.9*	12.2	1.4
Improves the chance of spontaneous delivery	74	14.9	52.7*	24.3	8.1	0.0
Reduces maternal motor block	74	40.5	51.4*	5.4	2.7	0.0
Reduces maternal side-effects	74	14.9	41.9*	25.7	17.6	0.0
Reduces fetal and neonatal side-effects	74	4.1	31.1	35.1*	28.4	1.4
15. Single-injection spinal opioids with or without local anesthetics versus parenteral opioids:						
Improve analgesia	74	68.9*	28.4	2.7	0.0	0.0
Increase the duration of labor	74	1.4	5.4	20.3	51.4*	21.6
Decrease the chance of spontaneous delivery	74	1.4	8.1	10.8	54.1*	25.7
Increase maternal side-effects	74	0.0	25.7	25.7*	36.5	12.2
Increase fetal and neonatal side-effects	74	0.0	9.5	16.2	51.4*	23.0
16. Single-injection spinal opioids with local anesthetics versus spinal opioids without local anesthetics:						
Improve analgesia	74	44.6	44.6*	4.1	5.4	1.4
Increase the duration of labor	74	2.7	6.8	25.7	51.4*	13.5
Decrease the chance of spontaneous delivery	74	2.7	5.4	23.0	58.1*	10.8
Increase maternal motor block	74	13.5	54.1*	9.5	21.6	1.4
Increase maternal side-effects	74	1.4	27.0	23.0*	40.5	8.1
Increase fetal and neonatal side-effects	74	0.0	4.1	23.0	58.1*	14.9
Combined Spinal-Epidural (CSE) Techniques :						
17. CSE local anesthetics with opioids versus epidural local anesthetics with opioids:						
Improve early analgesia	74	48.6	35.1*	5.4	10.8	0.0
Improve overall analgesia	74	18.9	31.1	23.0*	25.7	1.4
Decrease the duration of labor	74	4.1	18.9	47.3*	29.7	0.0
Decrease the chance of spontaneous delivery	73	0.0	2.7	19.2	61.6*	16.4
Reduce maternal motor block	74	5.4	37.8	24.3*	32.4	0.0
Increase maternal side-effects	74	0.0	18.9	24.3	54.1*	2.7
Increase fetal and neonatal side-effects	74	0.0	5.4	27.0	55.4*	12.2

	N	Percent Responding to Each Item				
		Strongly Agree	Agree	Uncertain	Disagree	Strongly Disagree
Patient-Controlled Epidural Analgesia (PCEA):						
18. PCEA versus continuous infusion epidurals:						
Improves analgesia	75	16.0	41.3*	26.7	12.0	4.0
Improves maternal satisfaction	75	41.3	46.7*	8.0	2.7	1.3
Reduces the need for anesthetic interventions	75	42.7	36.0*	10.7	9.3	1.3
Increases the chance of spontaneous delivery	74	4.1	13.5	45.9*	33.8	2.7
Reduces maternal motor block	75	9.3	38.7	24.0*	26.7	1.3
Decrease maternal side-effects	75	5.3	28.0	30.7*	34.7	1.3
19. PCEA with a background infusion versus PCEA without a background infusion:						
Improves analgesia	74	23.0	54.1*	16.2	6.8	0.0
Improves maternal satisfaction	74	24.3	43.2*	23.0	9.5	0.0
Reduces the need for anesthetic interventions	74	21.6	56.8*	12.2	9.5	0.0
Decreases the chance of spontaneous delivery	74	0.0	4.1	41.9	51.4*	2.7
Increases maternal motor block	74	1.4	39.2	25.7*	32.4	1.4
Increases maternal side-effects	74	1.4	13.5	29.7	52.7*	2.7
Neuraxial Analgesia, Timing of Initiation, and Progress of Labor:						
20. Administering epidural analgesia at cervical dilations of < 5 centimeters (versus \geq 5 cm):						
Improves analgesia	75	50.7*	32.0	9.3	6.7	1.3
Reduces the duration of labor	75	0.0	6.7	45.3*	41.3	6.7
Improves the chance of spontaneous delivery	74	0.0	10.8	48.6*	32.4	8.1
Increases maternal motor block	75	1.3	28.0	17.3	42.7*	10.7
Increases maternal side-effects	75	1.3	5.3	20.0	61.3*	12.0
Increases fetal and neonatal side-effects	75	0.0	4.0	17.3	58.7*	20.0
21. Neuraxial techniques improve the likelihood of vaginal delivery for patients attempting vaginal birth after prior cesarean delivery						
	75	21.3	36.0*	33.3	8.0	1.3
Removal of Retained Placenta:						
22. If an epidural catheter is <i>in situ</i> and the patient is hemodynamically stable, epidural anesthesia is the preferred technique						
	75	66.7*	30.7	2.7	0.0	0.0
23. In cases involving major maternal hemorrhage, a general endotracheal anesthetic is preferred over neuraxial anesthesia						
	75	30.7	48.0*	12.0	6.7	2.9
24. Administration of nitroglycerin for uterine relaxation improves success at removing retained placenta						
	75	34.7	48.0*	9.3	6.7	1.3

	N	Percent Responding to Each Item				
		Strongly Agree	Agree	Uncertain	Disagree	Strongly Disagree
Anesthetic Choices for Cesarean Delivery:						
25. Equipment, facilities, and support personnel available in the labor and delivery operating suite should be comparable to that available in the main operating suite	74	82.4*	16.2	1.4	0.0	0.0
26. General anesthesia versus epidural anesthesia:						
Reduces time to skin incision	74	40.5	37.8*	8.1	9.5	4.1
Increases maternal complications	74	37.8	47.3*	9.5	5.4	0.0
Increases fetal and neonatal complications	74	14.9	28.4	24.3*	29.7	2.7
27. General anesthesia versus spinal anesthesia:						
Reduces time to skin incision	74	20.3	35.1*	12.2	28.4	4.1
Increases maternal complications	74	33.8	50.0*	6.8	8.1	1.4
Increases fetal and neonatal complications	74	12.2	28.4	23.0*	33.8	2.7
28. Epidural anesthesia versus spinal anesthesia:						
Increases time to skin incision	74	43.2	43.2*	8.1	5.4	0.0
Reduces quality of anesthesia	74	12.2	56.8*	9.5	17.6	4.1
Increases maternal complications	74	1.4	13.5	28.4	48.6*	8.1
29. CSE anesthesia versus epidural anesthesia:						
Improves anesthesia	73	20.5	47.9*	20.5	11.0	0.0
Reduces time to skin incision	73	17.8	53.4*	12.3	16.4	0.0
Reduces maternal side-effects	73	2.7	12.3	30.1	52.1*	2.7
30. CSE anesthesia versus spinal anesthesia:						
Improves anesthesia	72	1.4	15.3	25.0	52.8*	5.6
Increases flexibility for prolonged procedures	73	61.6*	32.9	4.1	1.4	0.0
Increases time to skin incision	73	6.8	49.3*	17.8	21.9	4.1
Reduces maternal side-effects	73	1.4	11.0	37.0	47.9*	2.9
31. Use of pencil-point spinal needles versus cutting-bevel spinal needles reduces maternal complications	73	75.3*	23.3	1.4	0.0	0.0
32. Intravenous fluid preloading versus no intravenous fluid preloading for spinal anesthesia reduces maternal hypotension	73	30.1	46.6*	12.3	9.6	1.4
33a. Intravenous ephedrine is an acceptable agent to treat hypotension during neuraxial anesthesia	75	48.0	49.3*	1.3	1.3	0.0
33b. Intravenous phenylephrine is an acceptable agent to treat hypotension during neuraxial anesthesia	75	50.7*	40.0	6.7	2.7	0.0

	N	<u>Percent Responding to Each Item</u>				
		<u>Strongly Agree</u>	<u>Agree</u>	<u>Uncertain</u>	<u>Disagree</u>	<u>Strongly Disagree</u>
34. Neuraxial opioids versus parenteral opioids for postoperative analgesia after regional anesthesia for cesarean delivery:						
Improves analgesia	69	60.9*	33.3	5.8	0.0	0.0
Improves maternal satisfaction	69	52.2*	33.3	8.7	5.8	0.0
Postpartum Tubal Ligation:						
35. Neuraxial versus general anesthesia reduces maternal complications	70	24.3	58.6*	12.9	2.9	1.4
36. An immediate (≤ 8 hours) postpartum tubal ligation does not increase maternal complications	70	14.3	50.0*	22.9	11.4	1.4
Management of Complications:						
37. Availability of resources for management of hemorrhagic emergencies reduces maternal complications	70	74.3*	25.7	0.0	0.0	0.0
38. Immediate availability of equipment for management of airway emergencies reduces maternal, fetal and neonatal complications	70	80.0*	20.0	0.0	0.0	0.0
39. Immediate availability of basic and advanced life-support equipment in the labor and delivery suite reduces maternal, fetal and neonatal complications	70	78.6*	21.4	0.0	0.0	0.0
40. Routine use of central venous or pulmonary artery catheterization reduces maternal complications in severe preeclamptic patients	70	0.0	10.0	12.9	55.7*	21.4

Table 6. ASA Membership Survey Responses ^{§§§}

	N	Percent Responding to Each Item				Strongly Disagree
		Strongly Agree	Agree	Uncertain	Disagree	
Perianesthetic Evaluation:						
1. Directed history and physical exam reduces maternal, fetal and neonatal complications	2324	57.5*	38.3	3.0	1.0	0.1
2. Communication between anesthetic and obstetric providers reduces maternal, fetal and neonatal complications	2321	77.9*	21.3	0.6	0.2	0.1
3. A routine intrapartum platelet count does not reduce maternal anesthetic complications	2320	11.9	36.2	22.3*	23.6	6.0
4. An intrapartum platelet count reduces maternal anesthetic complications:						
For suspected preeclampsia	2326	35.8	47.9*	11.4	4.3	0.6
For suspected coagulopathy	2323	46.8	43.5*	6.2	2.8	0.6
5. All parturients should have an intrapartum blood sample sent to the blood bank to reduce maternal complications	2317	22.1	34.3*	19.0	21.9	2.7
6. Perianesthetic recording of the fetal heart rate reduces fetal and neonatal complications	2319	25.0	38.5*	25.2	9.9	1.6
Aspiration Prophylaxis:						
7a. Oral intake of clear liquids <i>during</i> labor improves patient comfort and satisfaction	2283	15.4	65.5*	12.1	6.2	0.8
7b. Oral intake of clear liquids <i>during labor does not</i> increase maternal complications	2285	6.7	40.2	23.6*	23.5	6.0
8a. Oral intake of solids <i>during labor</i> increases maternal complications	2284	48.2	38.0*	9.9	2.8	1.1
8b. The patient undergoing elective cesarean delivery should undergo a fasting period for solids of 6 to 8 hours depending on the type of food ingested (e.g., fat content)	2283	66.8*	30.3	1.1	1.3	0.5
8c. The patient undergoing elective postpartum tubal ligation should undergo a fasting period for solids of 6 to 8 hours depending on the type of food ingested (e.g., fat content)	2281	66.9*	30.2	1.1	1.4	0.4
9. Administration of a nonparticulate antacid prior to operative procedures reduces maternal complications	2281	24.5	43.3*	24.0	7.2	1.1
Anesthetic Care for Labor and Delivery:						
 Neuraxial Techniques:						
10. Prophylactic spinal or epidural catheter insertion for complicated parturients reduces maternal complications	2071	17.6	42.4*	26.9	11.8	1.2

^{§§§} N = the number of members who responded to each item. An asterisk beside a percentage score indicates the median.

	N	Percent Responding to Each Item					Strongly Disagree
		Strongly Agree	Agree	Uncertain	Disagree		
11. Continuous epidural infusion using local anesthetics with or without opioids versus parenteral opioids:							
Improves analgesia	2170	73.6*	25.1	0.8	0.4	0.1	
Increases the duration of labor	2174	1.2	14.4	19.0	51.7*	13.8	
Decreases the chance of spontaneous delivery	2171	0.8	7.4	16.9	53.3*	21.6	
Increases maternal side-effects	2169	0.6	12.0	9.8	58.9*	18.7	
Increases fetal and neonatal side-effects	2168	0.3	3.0	7.5	61.3*	27.9	
12. Continuous epidural infusion using local anesthetics with or without opioids versus spinal opioids with or without local anesthetics:							
Improves analgesia	2160	17.4	36.5*	24.8	20.2	1.2	
Increases the duration of labor	2161	0.8	8.9	31.8	49.7*	8.8	
Decreases the chance of spontaneous delivery	2158	0.6	5.8	27.7	53.7*	12.3	
Increases maternal motor block	2149	3.7	36.0	16.1*	38.7	5.4	
Increases maternal side-effects	2152	0.7	10.2	21.9	58.4*	8.8	
Increases fetal and neonatal side-effects	2153	0.4	4.2	20.9	61.2*	13.3	
13a. Induction of epidural analgesia using local anesthetics with opioids versus epidural analgesia with equal concentrations of local anesthetics without opioids:							
Improves analgesia	2153	34.6	46.1*	6.2	10.8	2.3	
Increases maternal side-effects	2150	2.6	38.0	12.8*	40.4	6.2	
Increases fetal and neonatal side-effects	2142	0.7	7.5	17.5	63.1*	11.3	
13b. Induction of epidural analgesia using low-dose local anesthetics with opioids versus higher concentrations of epidural local anesthetics without opioids:							
Improves analgesia	2155	13.1	31.7	26.9*	26.6	1.7	
Increases maternal side-effects	2154	1.1	13.8	15.8	55.7*	13.6	
Increases fetal and neonatal side-effects	2147	0.6	4.5	19.3	60.8*	14.8	
14a. Maintenance of epidural infusion of lower concentrations of local anesthetics with opioids versus higher concentrations of local anesthetics without opioids:							
Improves analgesia	1977	17.2	38.5*	24.0	19.2	1.0	
Reduces the duration of labor	1980	3.9	28.0	44.9*	21.6	1.6	
Improves the chance of spontaneous delivery	1977	6.9	41.1	35.9*	15.1	1.0	
Reduces maternal motor block	1977	31.3	63.0*	2.9	2.4	0.4	
Reduces maternal side-effects	1971	11.4	47.1*	26.8	14.0	0.9	
Reduces fetal and neonatal side-effects	1972	7.4	34.4	38.1*	18.6	1.5	

	N	Percent Responding to Each Item					Strongly Disagree
		Strongly Agree	Agree	Uncertain	Disagree		
14b. Maintenance of epidural analgesia using bupivacaine \leq 0.125% with opioids versus bupivacaine concentrations $>$ 0.125% without opioids:							
Improves analgesia	1973	16.5	38.6*	23.9	19.7	1.4	
Reduces the duration of labor	1975	4.4	25.6	46.9*	21.5	1.7	
Improves the chance of spontaneous delivery	1973	6.1	36.9	38.9*	16.7	1.4	
Reduces maternal motor block	1967	23.4	63.7*	5.3	6.5	1.1	
Reduces maternal side-effects	1960	9.2	44.7*	27.0	18.1	1.0	
Reduces fetal and neonatal side-effects	1957	6.3	31.3	39.0*	21.6	1.8	
15. Single-injection spinal opioids with or without local anesthetics versus parenteral opioids:							
Improve analgesia	1966	36.9	50.2*	8.9	3.6	0.5	
Increase the duration of labor	1963	0.4	2.7	31.5	55.8*	9.6	
Decrease the chance of spontaneous delivery	1967	0.4	2.8	27.9	58.3*	10.7	
Increase maternal side-effects	1958	2.1	23.7	23.1	45.1*	5.8	
Increase fetal and neonatal side-effects	1960	0.7	7.7	25.6	55.9*	10.2	
16. Single-injection spinal opioids with local anesthetics versus spinal opioids without local anesthetics:							
Improve analgesia	1961	29.2	55.6*	9.4	5.5	0.4	
Increase the duration of labor	1960	1.1	10.2	43.0*	41.2	4.6	
Decrease the chance of spontaneous delivery	1959	0.8	8.1	38.4	47.1*	5.7	
Increase maternal motor block	1955	12.5	59.0*	11.6	15.4	1.4	
Increase maternal side-effects	1951	2.5	33.1	28.9*	33.1	2.4	
Increase fetal and neonatal side-effects	1954	1.0	11.3	36.2	46.8*	4.7	
Combined Spinal-Epidural (CSE) Techniques :							
17. CSE local anesthetics with opioids versus epidural local anesthetics with opioids:							
Improve early analgesia	1887	31.1	44.6*	11.7	11.2	1.5	
Improve overall analgesia	1884	14.0	26.8	27.1*	28.7	3.5	
Decrease the duration of labor	1884	1.5	8.8	48.2*	38.2	3.4	
Decrease the chance of spontaneous delivery	1882	0.3	3.1	38.5	52.1*	6.0	
Reduce maternal motor block	1880	4.0	23.8	27.6*	41.0	3.5	
Increase maternal side-effects	1877	2.0	28.2	33.0*	34.2	2.6	
Increase fetal and neonatal side-effects	1872	0.9	11.4	37.1	45.3*	5.2	

	N	Percent Responding to Each Item					Strongly Disagree
		Strongly Agree	Agree	Uncertain	Disagree		
Patient-Controlled Epidural Analgesia (PCEA):							
18. PCEA versus continuous infusion epidurals:							
Improves analgesia	1852	15.3	40.1*	29.2	14.6	0.8	
Improves maternal satisfaction	1848	27.8	46.5*	19.6	5.6	0.5	
Reduces the need for anesthetic interventions	1849	22.4	42.9*	21.4	12.1	1.1	
Increases the chance of spontaneous delivery	1845	2.6	12.1	56.9*	26.4	2.1	
Reduces maternal motor block	1846	4.3	34.1	40.4*	20.5	0.8	
Decrease maternal side-effects	1838	3.8	27.0	46.5*	21.9	0.9	
19. PCEA with a background infusion versus PCEA without a background infusion:							
Improves analgesia	1840	26.0	48.4*	20.8	4.7	0.3	
Improves maternal satisfaction	1840	25.4	46.0*	24.1	4.2	0.3	
Reduces the need for anesthetic interventions	1829	22.4	46.0*	24.7	6.6	0.3	
Decreases the chance of spontaneous delivery	1831	0.8	4.3	48.6*	41.6	4.8	
Increases maternal motor block	1837	1.0	27.3	40.8*	28.6	2.2	
Increases maternal side-effects	1828	0.8	12.8	43.5*	39.6	3.3	
Neuraxial Analgesia, Timing of Initiation, and Progress of Labor:							
20. Administering epidural analgesia at cervical dilations of < 5 centimeters (versus \geq 5 cm):							
Improves analgesia	1831	25.9	52.7*	10.4	10.1	0.9	
Reduces the duration of labor	1825	1.9	13.5	40.1*	41.2	3.4	
Improves the chance of spontaneous delivery	1823	1.8	14.9	49.4*	30.9	3.0	
Increases maternal motor block	1819	0.9	20.5	21.2	53.4*	4.0	
Increases maternal side-effects	1821	0.7	11.0	22.3	61.1*	5.0	
Increases fetal and neonatal side-effects	1820	0.3	4.3	23.0	64.6*	7.7	
21. Neuraxial techniques improve the likelihood of vaginal delivery for patients attempting vaginal birth after prior cesarean delivery							
	1816	8.7	41.6*	37.9	10.1	1.7	
Removal of Retained Placenta:							
22. If an epidural catheter is <i>in situ</i> and the patient is hemodynamically stable, epidural anesthesia is the preferred technique							
	1821	30.8	59.5*	4.3	4.4	1.0	
23. In cases involving major maternal hemorrhage, a general endotracheal anesthetic is preferred over neuraxial anesthesia							
	1823	36.0	48.8*	6.9	7.5	0.9	
24. Administration of nitroglycerin for uterine relaxation improves success at removing retained placenta							
	1812	15.6	54.1*	26.4	3.5	0.4	

	N	Percent Responding to Each Item				
		Strongly Agree	Agree	Uncertain	Disagree	Strongly Disagree
Anesthetic Choices for Cesarean Delivery:						
25. Equipment, facilities, and support personnel available in the labor and delivery operating suite should be comparable to that available in the main operating suite	1815	78.3*	20.3	0.5	0.9	0.1
26. General anesthesia versus epidural anesthesia:						
Reduces time to skin incision	1826	30.9	46.3*	6.8	14.3	1.6
Increases maternal complications	1824	27.3	50.1*	10.9	9.6	2.0
Increases fetal and neonatal complications	1825	13.9	37.5*	23.2	22.8	2.6
27. General anesthesia versus spinal anesthesia:						
Reduces time to skin incision	1823	13.1	37.2*	13.7	30.1	6.0
Increases maternal complications	1815	23.8	49.6*	10.7	13.8	2.0
Increases fetal and neonatal complications	1803	13.6	37.2*	21.9	24.6	2.8
28. Epidural anesthesia versus spinal anesthesia:						
Increases time to skin incision	1823	32.1	54.3*	3.8	8.7	1.0
Reduces quality of anesthesia	1821	15.0	51.0*	8.8	21.5	3.6
Increases maternal complications	1816	1.2	8.8	24.5	59.1*	6.4
29. CSE anesthesia versus epidural anesthesia:						
Improves anesthesia	1794	18.7	45.4*	22.6	12.5	0.9
Reduces time to skin incision	1795	14.7	38.2*	21.7	23.2	2.3
Reduces maternal side-effects	1791	2.6	9.4	42.4*	43.3	2.4
30. CSE anesthesia versus spinal anesthesia:						
Improves anesthesia	1800	4.4	14.3	28.5	48.2*	4.6
Increases flexibility for prolonged procedures	1808	32.1	54.8*	10.2	2.5	0.4
Increases time to skin incision	1804	9.9	48.7*	17.7	22.1	1.7
Reduces maternal side-effects	1802	0.9	7.7	41.6*	46.1	3.7
31. Use of pencil-point spinal needles versus cutting-bevel spinal needles reduces maternal complications	1819	51.7*	39.4	5.7	2.9	0.4
32. Intravenous fluid preloading versus no intravenous fluid preloading for spinal anesthesia reduces maternal hypotension	1817	40.0	43.0*	9.0	6.5	1.4
33a. Intravenous ephedrine is an acceptable agent to treat hypotension during neuraxial anesthesia	1819	50.7*	47.3	0.9	1.0	0.1
33b. Intravenous phenylephrine is an acceptable agent to treat hypotension during neuraxial anesthesia	1820	31.9	52.8*	6.0	8.0	1.3

	N	<u>Percent Responding to Each Item</u>					<u>Strongly Disagree</u>
		<u>Strongly Agree</u>	<u>Agree</u>	<u>Uncertain</u>	<u>Disagree</u>		
34. Neuraxial opioids versus parenteral opioids for postoperative analgesia after regional anesthesia for cesarean delivery:							
Improves analgesia	1822	40.1	49.7*	6.9	3.0	0.3	
Improves maternal satisfaction	1816	35.0	47.4*	13.0	4.1	0.6	
Postpartum Tubal Ligation:							
35. Neuraxial versus general anesthesia reduces maternal complications	1812	28.8	45.0*	15.2	9.4	1.6	
36. An immediate (≤ 8 hours) postpartum tubal ligation does not increase maternal complications	1814	6.4	34.1	32.3*	23.0	4.2	
Management of Complications:							
37. Availability of resources for management of hemorrhagic emergencies reduces maternal complications	1823	67.9*	30.8	1.0	0.3	0.0	
38. Immediate availability of equipment for management of airway emergencies reduces maternal, fetal and neonatal complications	1817	77.2*	22.1	0.6	0.2	0.0	
39. Immediate availability of basic and advanced life-support equipment in the labor and delivery suite reduces maternal, fetal and neonatal complications	1812	73.4*	24.8	1.6	0.2	0.0	
40. Routine use of central venous or pulmonary artery catheterization reduces maternal complications in severe preeclamptic patients	1822	3.2	13.3	33.0	40.8*	9.6	